

EMBOTAPE

Pyrantel embonate 11.4g/syringe



BENEFITS

- For the treatment and control of adult infections of large and small redworms ascarids and pinworms.
- For the treatment and control of tapeworm
- Calibrated syringe for accurate dosage
- Short withdrawal 7 days
- Broad spectrum activity of pyrantel embonate

PACKAGING

| LIST NO. | UNIT PACKAGE | CASE SIZE |
|----------|---------------|-----------|
| 1EMB001 | 28.5G SYRINGE | 20 |

See reverse side for full indications, administration and dosage.



EMBOTAPE

Presentation

Embotape is a ready to administer oral wormer for horses and other equines. Each 28.5g syringe contains Pyrantel Embonate 11.4g as active ingredient and Butylated Hydroxy Toluene 0.0057g as an anti-oxidant. Each 100kg dose contains 1.9g Pyrantel Embonate .

Uses

For the treatment and control of adult infections of large and small Redworms, Ascarids and Pinworms found in the gastro-intestinal tract of horses and other equines. Embotape also treats and controls tapeworm infection.

Dosage and Administration

For direct oral administration only.
One syringe per 600kg bodyweight as a single dose (which is equivalent to one syringe mark per 100kg bodyweight).

Control and treatment of Strongyles, Oxyuris and Parascaris:

Embotape should be used at a dose rate of 19mg pyrantel embonate per kg bodyweight. The dosing programmes are as follows:

- A Foals (1 - 8 months of age): dose every 4 weeks.
- B Horses (over 8 months of age): dose every 6 - 8 weeks, but during the summer and autumn when at grass dose every 4 - 6 weeks. Always dose 3 - 4 days before turning out after in-wintering.
- C Suckler mares: It has been shown that reduction of strongyle challenge to the suckling foal at pasture can be achieved by using clean pasture (re-seeded or not grazed the previous year by horses), dosing the mare 3 - 4 days before turning out and then at intervals of 2 - 4 weeks until the end of Autumn. Ideally mares with foals should go out to clean pasture or, if this is not possible, delay turning them out until June.

The prescribed amount of Embotape is deposited on the tongue of the animal and the animal allowed to swallow. The complete content of one syringe contains 11.4g pyrantel embonate (6 graduated doses of 1.9g) in 28.5g paste and is sufficient for the treatment of 600kg bodyweight. Each graduation of the syringe is sufficient for the treatment of 100kg bodyweight.

| Weight | Type | Routine Treatment |
|----------------|--|---|
| Up to 100kg | Miniature ponies | Syringe mark 100kg |
| 101 - 300kg | Donkey, Shetland and other small ponies, foals | Syringe mark 300kg (½ syringe) |
| 301 - 400kg | Dartmoor, New Forest, Welsh | Syringe mark 400kg |
| 401 - 500kg | Light hunters, Arabs, etc. | Syringe mark 500kg |
| 501 - 600kg | Thoroughbreds | Syringe mark 600kg (1 syringe) |
| 601kg and over | Heavy hunters, draught horses | 1 full syringe plus additional 100kg syringe marks for each extra 100kg bodyweight. |

Control and treatment of anoplocephala (tapeworm):

Embotape should be used at a dose rate of 38mg pyrantel embonate per kg bodyweight (ie twice the dose used for strongyles). The need for retreatment may vary, but if considered necessary, should be carried out after an interval of 6 weeks.

Contra-indications, Warnings, etc.

The same syringe should only be used to dose two animals if they are both healthy and are either grazing together, or are on the same premises and in direct contact with each other. Impaction of the small intestine may occur in foals heavily infected with Ascarids. Colic symptoms may be seen as soon as 30 minutes after treatment.
Do not use in severely debilitated animals.
Do not use in horses with known sensitivity to the active ingredient.
In case of diminished efficacy consult a veterinarian.

Operator Warnings

Combined administration of pyrantel and levamisole or piperazine is not recommended.
Avoid contact with skin; wash hands and any other parts of the body which come into contact with the product after use. Avoid handling the product if you are known to be hypersensitive to pyrantel.

Withdrawal Periods

Horses may be slaughtered for human consumption only after 7 days from the last treatment.

Storage Precautions

Keep out of reach and sight of children.
Protect from direct sunlight.
On occasions where only part of the paste is used, replace the cap and store as indicated.

Legal Category LM
Package Quantities 28.5g syringe x 20.
VPA 10960/25/1

This information should be used as a general guide, for more specific instructions on the use of Bimeda[®] products always read the product label carefully.

