

# HYSPAN

*Neomycin Sulphate*



## BENEFITS

- A comprehensive oral scour treatment in calves
- Unique formulation to combat infection, loss of electrolytes and dehydration
- Contains the powerful antibiotic Neomycin sulphate, also contains glucose and electrolytes
- Excellent in the fight against scours caused by E-coli and salmonella
- Readily dispersable in warm water for quick uptake by animal after administration

## PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1HYS002	100g	20

See reverse side for full indications, administration and dosage.



# HYSPAN 400 mg ORAL POWDER

## Presentation

It is a white to off-white free-flowing powder for oral administration. Each 100g sachet contains 400mg of Neomycin Sulphate.

## Target Species

Calves.

## Indications for use, specifying the target species

For the treatment of scour in calves mainly due to *Escherichia coli* and *Salmonella*.  
Specially formulated to combat infection, dehydration and loss of electrolytes.

## Contraindications

Do not use in animals with known hypersensitivity to the active ingredients.

## Special warnings for each target species

Withdraw all milk and milk replacer from scouring animals.

## Special precautions for use

### Special precautions for use in animals

Withdraw all milk and milk replacer from scouring animals. Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

## Amounts to be administered and administration route

For oral administration.  
For calves between 35-75kg, 5 sachets are required. Administer the contents of 1 sachet dispersed in 3 pints of warm water twice daily for 2 days and once on the third day. Start treatment as soon as possible after the start of diarrhoea.

Discontinue treatment after 2 days if no response is observed and consult your veterinary surgeon. Do not extend treatment beyond the recommended time

## Withdrawal period(s)

Animals intended for human consumption should not be slaughtered during treatment. Animals intended for human consumption should not be slaughtered until 28 days after the last treatment.

## Pharmacodynamic properties

Hyspan acts locally on the gastrointestinal tract. Neomycin sulphate is an antibiotic which is poorly absorbed from the gut and is thus present in maximum concentration at the site of infection. The electrolytes restore electrolyte balance and relieve dehydration. Glucose monohydrate is an immediate source of energy which enhances the absorption of electrolytes and water thus speeding rehydration.

## Shelf life

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.

## Special precautions for storage

Store below 25°C.

## Nature and contents of immediate packaging

Foil laminate sachet containing 100g of product.

## Special precautions for the disposal of unused veterinary medicinal product or waste materials

Unused product or waste material should be disposed of in accordance with current practice for pharmaceutical waste under National Waste Disposal Regulations.

VPA

10960/067/001

This information should be used as a general guide, for more specific instructions on the use of Bimeda<sup>®</sup> products always read the product label carefully.

