

BILOSIN

200

Tylosin 200mg



BENEFITS

- Tylosin base injection
- Excellent syringability
- Effective against a wide range of Gram-negative and Gram-positive organisms

PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1BIL001	100ml	12

See reverse side for full indications, administration and dosage.



BILOSIN 200 INJECTION

Presentation

A clear yellow sterile solution suitable for parenteral administration. Each ml contains 200mg Tylosin Base BP(Vet). Also contains Benzyl Alcohol 4% as a preservative.

Uses

For the treatment of diseases in pigs involving organisms sensitive to Tylosin, including swine erysipelas (*Erysipelothrix rhusiopathiae*) vibronic dysentery and pneumonia (*Mycoplasma hyopneumoniae*).

Dosage and Administration

0.5ml/10kg bodyweight, equivalent to 10mg of Tylosin per kg bodyweight, by intramuscular injection every 12 hours, up to a maximum of 6 injections. Do not inject more than 5ml at a single injection site. If a larger injection volume is necessary, it should be divided and administered at different injection sites.

Contra-indications, Warnings, etc.

Do not use in animals known to be hypersensitive to the active ingredient.

Avoid skin contact with the preparation.
Wash hands after use.

Withdrawal Periods

Pigs must not be slaughtered for human consumption during treatment. Pigs may be slaughtered for human consumption only after 21 days from the last treatment.

Pharmaceutical Precautions

Do not store above 25°C.

Keep out of reach and sight of children.
Following withdrawal of the first dose, use the product within 28 days.
Avoid the introduction of contamination during use. Should any apparent growth or discoloration occur, the product should be discarded.

Legal Category POM

Package Quantities

Multi-dose vials of 100ml.

VPA 10126/40/1

This information should be used as a general guide, for more specific instructions on the use of Bimeda[®] products always read the product label carefully.

