

BIMOXYL L.A

150mg Amoxicillin/ml



BENEFITS

- Rapid absorption due to the patented penetration enhancer incorporated into the formulation
- Starts to work straight away with prolonged activity for 48 hours
- Well tolerated in cattle, sheep pigs and dogs.
- Broad spectrum antibiotic offering cover against Gram-negative and Gram-positive organisms
- Effective against diseases of the respiratory, alimentary and urogenital tracts
- Widespread tissue distribution
- Excellent syringeability via 21 gauge needle
- Single shot treatment in most cases

PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1BIM027	100ml	12

See reverse side for full indications, administration and dosage.



BIMOXYL LA INJECTION

Presentation

A sterile, off-white, oily, injectable suspension for intramuscular administration. Each ml contains 150mg Amoxycillin as Amoxycillin Trihydrate in a long acting oily base.

Uses

Cattle

For the control and treatment of respiratory and other infections caused by amoxycillin susceptible Gram-positive and Gram-negative bacteria only.

Sheep, Pigs and Dogs

For the treatment of infectious diseases in pigs, sheep and dogs, caused by or associated with organisms sensitive to amoxycillin.

Dosage and Administration

Cattle, Sheep and Pigs: By intramuscular route only.

Dogs: By subcutaneous injection.

The injection site should be massaged after injection.

The recommended dosage rate is 15mg amoxycillin per kg bodyweight. This is equivalent to 1ml/10kg. One repeat administration may be given after 48 hours.

The maximum injection volume at any one site is:

Cattle: 20ml; Sheep: 4ml; Pigs: 5ml; Dogs: 2.5ml.

Larger dose volumes should be divided and given into separate sites.

Use a dry sterile needle and syringe for extraction of suspension to avoid hydrolysis of amoxycillin.

Contra-indications, Warnings, etc.

For animal treatment only.

Not suitable for intravenous or intrathecal administration.

Not to be administered to small herbivores.

Not for use in ewes producing milk for human consumption or food processing.

Do not use in known cases of hypersensitivity to penicillin or the active ingredient.

Occasional local reaction of a transient nature may occur at the site of injection.

Bimoxyl LA is unlikely to interact significantly with any of the other drugs commonly administered to animals.

There is no evidence that the use of amoxycillin presents any particular hazard to the dam or to the foetus.

As with

all other antibiotics, Bimoxyl LA should be used with caution during pregnancy and lactation.

Operator Warnings

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

- 1 Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations.
- 2 Handle this product with great care to avoid exposure, taking all recommended precautions.
- 3 If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Withdrawal Periods

Animals must not be slaughtered for human consumption during treatment.

Cattle may be slaughtered for human consumption only after 18 days from the last treatment. Sheep may be slaughtered for human consumption only after 21 days from the last treatment. Pigs may be slaughtered for human consumption only after 11 days from the last treatment.

Not for use in ewes producing milk for human consumption or food processing.

Milk for human consumption must not be taken during treatment. Milk for human consumption may be taken only from cows from 72 hours (ie at the 6th milking) from the last treatment.

Pharmaceutical Precautions

Do not store above 25°C.

Keep out of reach and sight of children.

Shake well before use.

Swab the septum before removing each dose.

Use a dry, sterile needle and syringe.

Following withdrawal of the first dose, use the product within 28 days. Discard unused material.

Avoid the introduction of contamination during use. Should any apparent growth or discoloration occur, the product should be discarded.

Legal Category

POM

Package Quantities

Multidose vials of 100ml

VPA

10126/9/1

This information should be used as a general guide, for more specific instructions on the use of Bimeda® products always read the product label carefully.

