

# Oxycomplex NS

**Oxytetracycline hydrochloride**  
**Flunixin Meglumine**



## BENEFITS

- A powerful combination product with a proven antibiotic and a potent anti-inflammatory agent
- Broad spectrum activity against Gram-negative and Gram-positive bacteria
- Rapid response and dramatic results
- For treatment of pneumonia, mastitis, metritis & gastroenteritis

## PACKAGING

| LIST NO. | UNIT PACKAGE | CASE SIZE |
|----------|--------------|-----------|
| 1OXY003  | 100ml        | 12        |

See reverse side for full indications, administration and dosage.



## OXYCOMPLEX NS

### Presentation

A clear, sterile, yellow to amber multidose aqueous solution for parenteral administration to cattle.

Each ml contains:

|  |       |
|--|-------|
| Oxytetracycline<br>(as Oxytetracycline Hydrochlorider) | 100mg |
| Flunixin<br>(as Flunixin Meglumine)                    | 20mg  |

Also contains N-methylpyrrolidone as a co-solvent and Sodium Formaldehyde Sulfoxylate as an antioxidant.

### Uses

For the control and treatment of infectious diseases of cattle, caused by or associated with organisms sensitive to oxytetracycline, where concurrent analgesic, anti-inflammatory, anti-endotoxic or antipyretic therapy is desired.

Oxycomplex NS is especially indicated for the treatment of respiratory disease (particularly that associated with Pasteurella infection) and acute mastitis (in conjunction with appropriate intramammary therapy).

### Dosage and Administration

By intravenous or deep intramuscular injection at a rate of 1ml of Oxycomplex NS per 10kg bodyweight (equivalent to 10mg oxytetracycline and 2mg flunixin per kg bodyweight). Treatment may be repeated daily for up to 5 days. Do not inject more than 20ml intramuscularly at a single site. Where the dose exceeds 20ml it should be divided between two or more sites, as appropriate.

### Contra-indications, Warnings, etc.

For animal treatment only.  
Not to be used in animals known to be hypersensitive to any of the active ingredients.  
Do not use in horses or donkeys.  
Do not exceed the stated dose on duration of the treatment.  
Do not administer other NSAIDs concurrently or within 24 hours of each other.  
Do not use in animals suffering from cardiac, hepatic or renal disease, where there is the possibility of gastro-intestinal ulceration or bleeding, where there is evidence of a blood dyscrasia. Prolonged use of NSAIDs, including flunixin, may predispose or lead to gastrointestinal ulceration. This product may be used in pregnant and lactating animals. Oxycomplex NS may potentiate the effects of warfarin and related drugs.

Because of their common mode of action, flunixin may potentiate and be potentiated by other NSAIDs, which act by interfering with prostaglandin synthesis.

Where other products are to be administered concurrently with Oxycomplex NS, drug compatibility should be carefully monitored.

Corticosteroids should not be used concurrently with this product.

Overdosing by intramuscular injection may give rise to swelling at the site of injection. Treatment should be symptomatic.

### Withdrawal Periods

Animals intended for human consumption must not be slaughtered during treatment.

Cattle intended for human consumption may only be slaughtered from 28 days after the last treatment.

Milk intended for human consumption must not be taken during

treatment or for 10 milkings (5 days) thereafter. Where cows are milked twice daily, milk may be taken for human consumption from the 11th milking following the last treatment.

### Pharmaceutical Precautions

Keep out of reach and sight of children.

Do not store above 25°C.

Protect from light.

Following withdrawal of the first dose, use the product within 28 days.

Unused product or waste material should be disposed of in accordance with current practice for pharmaceutical waste under national waste disposal regulations.

Avoid the introduction of contamination during use.

Should any apparent growth occur, the product should be discarded.

**Legal Category** POM

### Package Quantities

100ml vials.

**VPA** 10126/63/1

**This information should be used as a general guide, for more specific instructions on the use of Bimeda® products always read the product label carefully.**

