

TETROXY LA

Oxytetracycline



BENEFITS

- Convenience, one treatment usually sufficient, reduces labor and number of injection sites
- Broad spectrum activity of oxytetracycline
- Minimal irritancy-well tolerated in cattle, sheep and pigs
- Ease of administration- 1ml/10kg with excellent syringeability
- Rapid initial blood levels followed by a long acting effect
- Ready to use, no mixing or refrigeration
- High syringability, injects well even at cold temperatures

PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1TET015	100ml	12

See reverse side for full indications, administration and dosage.



TETROXY LA

Presentation

A long acting oxytetracycline injectable solution.
Each ml contains: 200mg oxytetracycline (as Oxytetracycline Dihydrate). This product does not contain an antimicrobial preservative Also contains Povidone and NMP as co-solvents and Sodium Formaldehyde Sulfoxylate 0.27% as an antioxidant.

Uses

For the treatment and control of diseases in cattle, sheep and pigs caused by or associated with organisms sensitive to oxytetracycline. These include: Pasteurella spp., Salmonella spp. Escherichia coli, Listeria spp

Dosage and Administration

By deep intramuscular injection.
1ml per 10kg bodyweight, equivalent to 20mg oxytetracycline per kg bodyweight.

Maximum volume at one injection site:
Cattle, Sheep and Pigs - 10ml.
Pigs under 10kg, maximum dose 1ml.

Because of the sustained blood levels attained at the above dosage rates with Tetroxy LA one treatment is usually sufficient.

Contra-indications, Warnings, etc.

For animal treatment only.
Not to be used in cats, dogs, horses and donkeys.
Not for use in ewes producing milk for human consumption.
The use of Tetroxy LA during the period of tooth development including late pregnancy, may lead to tooth discoloration. Occasionally local reaction of a transient nature may occur at the site of injection.

Operator Warnings

Avoid contact with the eyes.
Wash hands after use.

Withdrawal Periods

Cattle and sheep for human consumption should not be slaughtered within 28 days of treatment.

Pigs for human consumption should not be slaughtered within 21 days of treatment.
Not for use in sheep producing milk for human consumption.

Milk intended for human consumption or the manufacture of cheese and yoghurt should not be drawn from treated animals within 7 days after treatment. Milk may only be taken from the 15th milking with a twice a day milking programme.

Pharmaceutical Precautions

Keep out of reach and sight of children.
Do not store above 25°C.
Avoid the introduction of contamination during use.
Should any apparent growth occur, the product should be discarded.
Following withdrawal of the first dose, the product should be used within 28 days.

The product should not be diluted with solutions of calcium salts as this causes precipitation.
Unused product or waste material should be disposed of in accordance with current practice for pharmaceutical waste under national waste disposal regulations.

Legal Category **POM**

Package Quantities

100ml multidose injection

VPA **10126/10/1**

This information should be used as a general guide, for more specific instructions on the use of Bimeda[®] products always read the product label carefully.

