


TETROXY EGG FORMULA

INDICATIONS

A broad-spectrum antibiotic with essential vitamins. For use in drinking water to improve growth rate and food conversion in poultry and to improve and maintain egg production.

BENEFITS

- Provides a convenient method of adding antibiotic and vitamins to the drinking water or rations.
- · Prolongs the laying period.
- Improves feed conversion and increases egg numbers.
- In breeding stock, it increases egg production, improves egg fertility and increases the number of hatched chicks.



PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1TET073	120g sachet	4

See reverse side for full indications, administration and dosage.



Bimeda®
Broomhill Road
Tallaght
Dublin 24, Ireland

Tel: 1850-515253 Fax: 01 451 5803

TECHNICAL INFORMATION



TETROXY EGG FORMULA

PRESENTATION

Tetroxy Egg is a water-soluble powder for oral administration. This medicated feed additive is indicated for use in laying birds.

INDICATIONS

A broad-spectrum antibiotic with essential vitamins. For use in drinking water to improve growth rate and food conversion in poultry and to improve and maintain egg production.

SUGGESTED DAILY DOSE

For more early eggs:

When laying starts add 5g to each 10 litres of the drinking water use at this level for 4 weeks. Thereafter, add 5g to each 50 litres of the drinking water.

To maintain improved egg production:

During the normal laying period add 5g to 50 litres of drinking water routinely.

During Stress Periods:

Or when disease is present increase dose to 5g in 5 litres of drinking water. Use until laying improves.

If production suddenly decreases:

Add 5g to 50 litres of drinking water until egg production returns to normal.

To prolong the laying period:

Use at the rate of 5g to 10 litres of water towards the end of the laying period.

For improved fertility:

In chicken and turkey breeding flocks use continuously at the rate of 5g to 10 litres of drinking water.

GUIDE TO DILUTION RATES:				
	NUMBER OF LITRES OF WATER TO BE TREATED			
	Under normal conditions	When laying starts	During period of stress and disease	
Quantity of Tetroxy Egg Formula required	5 g per 50 litres	5 g per 10 litres	5 g per 5 litres	
5 g (1 Teaspoonful)	50 litres	10 litres	5 litres	
50 g (10 Teaspoonful)	500 litres	100 litres	50 litres	
100 g	1000 litres	200 litres	100 litres	
200 g	2000 litres	400 litres	200 litres	

EACH 200g CONTAINS:

Active Substance

Oxytetracycline Hydrochloride	11g
Vitamin A	440,000 i.u.
Vitamin D3	79,100 i.u.
Vitamin E	132 i.u.
Vitamin B12	460 mcg
Vitamin K	186 mg
Nicotinamide	2,640 mg
Riboflavin	572 mg
Pantothenic Acid	927 mg
Carophyll Orange	800 mcg

WARNING

Poultry should not be slaughtered for human consumption during treatment.

For animal treatment only. Keep out of the reach of children

STORAGE CONDITIONS

Store in a cool, dry place.